


Contents lists available at [SciVerse ScienceDirect](http://SciVerse.ScienceDirect.com)

Pharmacology, Biochemistry and Behavior

journal homepage: www.elsevier.com/locate/pharmbiochembeh

Inhibition of nitric oxide synthase accentuates endotoxin-induced sickness behavior in mice

Deidiane E. Ribeiro^a, Viviane M. Maiolini^a, Roseli Soncini^a, José Antunes-Rodrigues^b, Lucila L.K. Elias^b, Fabiana C. Vilela^a, Alexandre Giusti-Paiva^{a,*}^a Department of Physiological Science, Institute of Biomedical Sciences, Federal University of Alfenas-MG, Brazil^b Department of Physiology, School of Medicine of Ribeirão Preto, University of São Paulo, Brazil

ARTICLE INFO

Article history:

Received 4 June 2012

Received in revised form 18 September 2012

Accepted 29 September 2012

Available online 6 October 2012

Keywords:

Endotoxin

Nitric oxide

Sepsis

Sickness behavior

ABSTRACT

Sickness behavior appears to be the expression of a central motivational state that reorganizes an organism's priorities to cope with infectious pathogens. To evaluate the possible participation of nitric oxide (NO) in lipopolysaccharide-induced sickness behaviors, mice were submitted to the forced swim test (FST), open field test and dark–light box test. Food intake and corticosterone plasma levels were evaluated. Lipopolysaccharide (LPS, 100 µg/kg; i.p.) administration increased the time spent floating in the FST and decreased locomotor activity in the open field. Indeed, treatment with LPS decreased the total number of transitions between the dark and light compartments of the apparatus. In addition, LPS reduced food intake and increased corticosterone levels. Pretreatment with L-NAME (30 mg/kg; i.p.) or aminoguanidine (50 mg/kg; i.p.) accentuated the behavioral changes induced by LPS in the FST, open field and light–dark box tests as well as induced an increment in hypophagia and in corticosterone levels. These findings confirm previous observations that have reported LPS-induced sickness behaviors. In addition, they provide evidence that the synthesis of NO modulates changes in depressive-like and exploratory behaviors in mice, which is supported by the fact that NO synthase inhibitors also attenuate LPS-induced behavioral changes. In addition, the present study suggests that NO may have a protective role, acting in an inhibitory feedback manner to limit LPS-induced sickness behavior.

© 2012 Elsevier Inc. Open access under the [Elsevier OA license](http://creativecommons.org/licenses/by/3.0/).

1. Introduction

Sickness behavior is an expression of a motivational state triggered by activation of the peripheral innate immune system whereby an organism reorganizes its priorities to fight infection (Parnet et al., 2002; Dantzer et al., 2008; Dantzer, 2009). Sickness behavior is easily induced in laboratory animals by the administration of endotoxins, which are the main active components of the outer membrane of gram-negative bacteria. Peripheral administration of endotoxin activates the immune system, resulting in the release of pro-inflammatory cytokines and prostaglandins (Long et al., 1990; Kelley et al., 2003; Engblom et al., 2002; Rorato et al., 2009; de Paiva et al., 2010). These peripherally produced cytokines gain access to the brain and produce the classical symptoms of sickness behavior, including reduction in locomotor activity and exploratory behaviors, anorexia and anhedonia (Engblom et al., 2002; Frenois et al., 2007). In addition, it has been demonstrated that upon stimulation by endotoxin and cytokines, nitric oxide synthase (NOS)

becomes expressed, leading to the production of large amounts of nitric oxide (NO) in peripheral tissues and in the brain (Moncada et al., 1991; Giusti-Paiva et al., 2005; Mollace et al., 2005).

NO is produced from L-arginine by three different isoforms of NOS, two of which are expressed constitutively, endothelial (eNOS) and neuronal (nNOS) (Mollace et al., 2005). Endotoxin induces the expression of another calcium-independent inducible isoform of NOS (iNOS) in a number of tissues including the brain (Moncada et al., 1991; Mollace et al., 2005; Cauwels, 2007). Pharmacological inhibition of NOS can be used to elucidate the contribution of NO, and previous reports support both a detrimental and beneficial role of NO during immune challenges. An increase in the amount of NO produced by iNOS might play a pivotal role in the pathophysiology of sepsis (Thiemermann, 1997; Szabo, 1996; Vincent et al., 2000; Cauwels, 2007).

Although a significant volume of literature indicates that cytokines and prostaglandins are important mediators of sickness behavior (Dunn and Swiergiel, 2005; Dantzer et al., 2008; de Paiva et al., 2010), there is a lack of data on the role of NO in endotoxin-induced sickness behavior. To evaluate the possible participation of NO in LPS-induced sickness behaviors, mice were submitted to well-accepted tests to evaluate depressive-like and exploratory behaviors, including the forced swim test (FST), open field test and dark–light box test.

* Corresponding author at: Institute of Biomedical Sciences, Federal University of Alfenas, MG, Rua Gabriel Monteiro da Silva, 700, 37130-000, Alfenas, MG, Brazil. Tel.: +55 35 3299 1303; fax: +55 35 3299 1063.

E-mail address: agiustipaiva@gmail.com (A. Giusti-Paiva).

2. Materials and methods

2.1. Animals

Adult male Swiss mice (22–30 g) were obtained from the Central Animal Facility of the Federal University of Alfenas. Animals were housed under controlled light (12:12 h light–dark cycle; lights on at 06:00 am) and temperature conditions (23 ± 1 °C) with access to water and food *ad libitum*. Animals were allowed to habituate to the housing facilities for at least 1 week before the experiments began. Behavioral studies were conducted in a quiet room between 09:00 and 11:00 a.m. to avoid circadian variation. All experiments were conducted in accordance with the Declaration of Helsinki addressing the welfare of experimental animals and with the approval of the Ethics Committee of the Federal University of Alfenas (protocol #0178/2008).

2.2. Experimental procedures

In the animal room, the mice were pretreated with non-selective NOS inhibitor (N ω -L-nitro-arginine methyl ester; L-NAME, 30 mg/kg), or a relatively selective iNOS inhibitor (aminoguanidine, 50 mg/kg), or vehicle (0.9% NaCl) 30 min before injections of lipopolysaccharide (LPS) serotype 026:B6 (100 μ g/kg; i.p.) or saline (0.9% NaCl). The behavioral tests were performed 120 min after the LPS treatment. These time points were chosen on the basis of previous behavioral, endocrine and neurochemical studies (Dunn and Swiergiel, 2005; Rorato et al., 2009; de Paiva et al., 2010). The doses of L-NAME and aminoguanidine used in the present study are in agreement with the doses commonly used in other reports that injected these compounds peripherally (Giusti-Paiva et al., 2004). All drugs were purchased from Sigma-Aldrich Co. and dissolved in sterile isotonic saline.

2.2.1. Forced swim test

This test was performed according to the method developed by Porsolt et al. for mice. Mice ($n = 12$ per group) were placed in a vertical glass cylinder (26 cm high, 12 cm diameter) filled with 25 °C water to a depth of 16 cm. Water depth was chosen to ensure that animals swam or floated without their hind limbs or tail touching the bottom. Each mouse was placed in the cylinder for 6 min, and the duration of floating (i.e., the time during which mice made only the smallest movements necessary to keep their heads above water) was scored. As suggested by Porsolt et al. (1977), only the data scored during the last 4 min were analyzed and presented (Porsolt et al., 1977; Dunn and Swiergiel, 2005; de Paiva et al., 2010).

2.2.2. Open field behavioral test

Locomotor activity was quantified for 5 min in an open field box, consisting of white Plexiglas 60 × 60 cm in diameter with a floor divided into 16 squares. Previous studies have indicated that this time-period was sufficient to produce differences between treatment groups. Furthermore, after 5 min, the mice habituate to the apparatus, thereby decreasing the differences between groups. Four squares were defined as the center and the 12 squares along the walls were considered the periphery. Each mouse ($n = 10$ per group) was gently placed in the exact center of the box. Activity was scored as a line crossing when a mouse removed all four paws from one square and entered another. Line crossings among the central four squares or among the peripheral 12 squares of the open field were counted separately (Dunn and Swiergiel, 2005; de Paiva et al., 2010).

2.2.3. Light–dark box test

The apparatus consisted of a Plexiglas rectangular box (48 cm long × 24 cm wide × 24 cm high) divided into a dark region (24 cm long) and a light region (24 cm long). The light and dark regions were separated by an opening (8.0 × 8.0 cm) that allowed the animals

to move between the two compartments. The dark region was made of black Plexiglas and covered with a black lid. The light portion was made of white Plexiglas, and a 60 W light was positioned directly over it. On the day of testing, each mouse was transported individually from the housing room to the testing room. Each mouse ($n = 10$ per group) was placed in the light compartment and allowed to move freely between the two compartments. The behavior was video-recorded for a total of 5 min, and the videotapes were scored for latency to the first transition and the number of transitions between the light and dark compartments (Lacosta et al., 1999; de Paiva et al., 2010).

2.2.4. Feeding behavior

The animals ($n = 10$ per group) fasted for 12 h before receiving injections. Immediately after injections, a fresh supply of pre-weighed food was given. Food intake was calculated at 2, 4, 6 and 24 h after the injection by measuring the difference between the pre-weighed standard chow available and the weight of chow and spilled crumbs at each time point. Changes in body weight were measured by weighing the animals at the beginning of the experiment as well as before and after an experimental day (Rorato et al., 2009; Soncini et al., 2012).

2.2.5. Plasma corticosterone assay

Trunk blood was collected from animals 2 h after LPS or saline injections in chilled heparinized tubes and was centrifuged at 3000 rpm for 15 min at 4 °C. Plasma was collected and stored at -20 °C. Plasma corticosterone was measured by radioimmunoassay.

2.2.6. Statistical analysis

The data obtained were analyzed using the GraphPad software program Version 4.0 and expressed as the mean \pm S.E.M. Statistically significant differences among groups were calculated by the application of an analysis of variance (ANOVA) followed by the Newman–Keuls test. P -values less than 0.05 ($p < 0.05$) were considered significant.




Fig. 1. Effects of pretreatment with vehicle, L-NAME (30 mg/kg) or aminoguanidine (AG, 50 mg/kg) on time spent floating in the forced swim test ($n = 12$ animals per group), which were measured 2 h after administration of either lipopolysaccharide (LPS) or saline. Each column represents the mean with S.E.M. The symbols denote significance levels: ** $p < 0.01$; *** $p < 0.001$ when compared with the control groups; # $p < 0.05$ when compared with the vehicle plus LPS group.




Fig. 2. Effects of pretreatment with vehicle, L-NAME (30 mg/kg) or aminoguanidine (AG, 50 mg/kg) on peripheral (A), central (B) and total (C) line crossings in the open field test ($n = 10$ per group), which were measured 2 h after administration of either LPS or saline. Each column represents the mean with S.E.M. The symbols denote significance levels: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$ when compared with the control groups; # $p < 0.05$; ## $p < 0.01$ when compared with the vehicle plus LPS group.

3. Results

3.1. Activity in the forced swim test

L-NAME and aminoguanidine did not alter the immobility period in the FST in saline treated mice. There was a significant increase in the immobility period 120 min after administration of LPS in the FST ($F_{5,71} = 16.94$; $p < 0.01$; Fig. 1) in mice pretreated with vehicle. The administration of LPS in mice previously treated with L-NAME or aminoguanidine caused a significantly higher immobility period in FST ($p < 0.05$), in comparison with the vehicle plus LPS group.

3.2. Activity in the open field test

Fig. 2 shows the effect of LPS on number of line crossings in the open field, and the modulatory effects of NOS inhibition. Pretreatment with L-NAME or aminoguanidine did not significantly change the locomotor activity in the open field 120 min after i.p. saline injection. On the other hand, LPS significantly decreased the number of line crossings in the center ($F_{5,59} = 11.9$, $p < 0.01$; Fig. 2A) and in the periphery ($F_{5,59} = 16.3$, $p < 0.01$; Fig. 2B), as well as the total number of line crossings ($F_{5,59} = 20.28$, $p < 0.001$; Fig. 2C). The number of line crossings in periphery and center of open field was even smaller in the mice pretreated with NOS inhibitors ($p < 0.05$).

3.3. Behavior in the light–dark box test

The amount of time spent before the first transition in the black–white box as well as the number of entries into the light compartment were evaluated (Fig. 3). There were no significant differences in latency to the first transition after LPS administration (Fig. 3A). However, there was a significant difference in the number of transitions between the light and dark compartments ($F_{5,59} = 28.8$; $p < 0.001$; Fig. 3B). The administration of LPS in mice previously treated with L-NAME or aminoguanidine caused a significant increase in latency to the first transition ($F_{5,59} = 4.01$; $p = 0.0036$; Fig. 3A), and an accentuated decrease in the number of transitions between the compartments, compared with the vehicle plus LPS group (Fig. 3B; $p < 0.05$).

3.4. Food intake and body weight

LPS significantly decreased food intake 4 h after injection ($F_{7,79} = 3.65$; $p < 0.01$; Fig. 4A), overnight food intake ($F_{7,79} = 6.466$; $p < 0.001$; Fig. 4A) and body weight ($F_{7,79} = 3.842$; $p < 0.01$; Fig. 4B). Furthermore, pretreatment with L-NAME or aminoguanidine significantly accentuated both hypophagic effects ($p < 0.05$) and reduced body weight gain ($p < 0.05$) induced by LPS.

3.5. Corticosterone plasma levels

Pretreatment with L-NAME or aminoguanidine did not significantly change the corticosterone plasma levels 2 h after i.p. saline injection; however, LPS injections resulted in an increase in the corticosterone plasma levels at this time. The administration of LPS in mice previously treated with L-NAME or aminoguanidine caused a significant increase in corticosterone secretion, compared with the control group (Fig. 5).

4. Discussion

The present study confirms previous observations that LPS can induce sickness behavior. These findings also provide evidence that NO acts as modulator for the development of depressive-like and exploratory behaviors in mice because non-selective NOS inhibition or selective inhibition of iNOS accentuated LPS-induced the behavioral changes.




Fig. 3. Effects of pretreatment with vehicle, L-NAME (30 mg/kg) or aminoguanidine (AG, 50 mg/kg) on latency to the first transition (A) and number of transitions (B) in the light–dark box test ($n = 10$ per group), which were measured 2 h after administration of either LPS or saline. Each column represents the mean with S.E.M. The symbols denote the significance levels: $^*p < 0.05$; $^{***}p < 0.001$ when compared with the control groups; $^{\#}p < 0.05$ when compared with the vehicle plus LPS group.

In these experiments, LPS administration increased the time spent floating in the FST and depressed the locomotor activity in the open field. These effects are characteristic of treatments that have been shown to induce depression-like behavior, and an accompanying deficit in locomotor activity would most likely affect performance in the FTS (Dunn and Swiergiel, 2005; de Paiva et al., 2010). In addition to promoting illness, LPS elicited an anxiogenic-like response in the light–dark box test. Indeed, treatment with LPS decreased the total number of transitions between dark and light compartments of the apparatus, which is a pattern believed to reflect anxiety. The proposition that LPS elicits anxiety is consistent with these findings that LPS induced an increase in corticotrophin-releasing factor (CRF) secretion, a hormone known to promote anxiety and activation of hypothalamus-pituitary-adrenal axis, as evidenced by increased corticosterone levels (Turnbull and Rivier, 1999; Rorato et al., 2009; Benedetti et al., 2011).

Activation of the immune system in response to LPS produced neural, neuroendocrine, and behavioral effects. These effects are mediated by circulating cytokines that influence brain activity by inducing the expression of neuronal NOS (nNOS) and inducible NOS (iNOS)

in the hypothalamus, and the expression of cyclooxygenase (COX)-2 and microsomal prostaglandin E synthase-1 (mPGES-1) in brain vascular cells (Engblom et al., 2002, 2003; Parnet et al., 2002). Previous work has indicated that fever is mediated by increased production of prostaglandin E_2 (PGE_2) that is synthesized along the blood–brain barrier (Engblom et al., 2003), which activates a population of target neurons in the thermoregulatory center of the brain with specific PGE_2 receptors (Ushikubi et al., 1998). In addition, sickness behavior in response to LPS has been shown to be absent or greatly attenuated in mice pre-treated with inhibitors of cyclooxygenase, thereby suggesting that sickness behavior induced by LPS is mainly prostaglandin dependent and is most likely mediated by a mechanism similar to febrile response mechanisms (Ushikubi et al., 1998; Dallaporta et al., 2007; de Paiva et al., 2010). At the behavioral level, sickness behavior is a motivational state that reorganizes an organism's priorities to cope with infectious pathogens; however, the mechanisms underlying sickness behaviors have not been fully elucidated (Dantzer, 2009; Kelley et al., 2003).

LPS is a potent inducer of iNOS. Large amounts of NO and peroxynitrite, among other factors, are responsible for the hypotension, vasoplegia, apoptosis, lactic acidosis and multi-organ failure observed in the late phases of endotoxic shock. The data supporting both a detrimental and beneficial role of NO in septic shock have changed the scientific impression of this molecule. Indeed, experimental and clinical use of NOS inhibitors, which do not differentiate clearly between constitutive endothelial NOS and iNOS, prevents LPS-induced hypotension (Giusti-Paiva et al., 2002, 2004; Vona-Davis et al., 2002). However, many detrimental effects of NOS inhibitors are also reported, including increases in pulmonary resistance, decreases in cardiac output and organ perfusion, and even an increase in mortality in experimental animals (Boyle et al., 2000; Metcalf et al., 2002; Vona-Davis et al., 2002). For this reason, selective iNOS inhibitors, such as aminoguanidine or thiourea derivatives, might be preferred over nonselective NOS inhibitors for the treatment of septic shock (Cauwels, 2007). However, because iNOS-derived NO is involved in multiple pathways, selective iNOS inhibition may be not as beneficial as expected. On the other hand, previous studies have demonstrated a relation between functional protection of inhaled NO and reduced leukocyte adhesion and emigration in inflammatory lung injury (Guidot et al., 1995; Trzeciak et al., 2008). Concurrently, we found that the preservation of oxygenation in animals given NO inhalation was associated with less pulmonary sequestration of leukocytes. Accordingly, inhalation of NO gas or NO-donors in septic shock might be a complementary treatment to the use of NOS inhibitors.

In addition to the constitutive isoforms nNOS, iNOS expression has also been detected in the brain during immune challenge. iNOS is preferentially expressed in macrophages and is up-regulated in response to a variety of inflammatory stimuli; however, iNOS has been reported in the brain following endotoxin administration (Minc-Golomb et al., 1994; Harada et al., 1999). Harada et al. (1999) have demonstrated that LPS activates both nNOS and iNOS gene expression and activity in the hypothalamus, with a delayed time-course expression of iNOS compared with nNOS. Overall, there is a profound increase in NO production derived from both nNOS and iNOS in the brain after systemic inflammation or sepsis (Minc-Golomb et al., 1994; Harada et al., 1999). Despite an increase in NO production, including in the brain, during endotoxemia in mice and because NOS inhibitors crosses the blood–brain barrier (Uribe et al., 1999; Mahar Doan et al., 2000), does NO can play a role in the central mechanisms the involvement of sickness behavior induced by LPS.

Previous reports suggest that one of the roles of endogenous NO might be to restrain activation of the hypothalamic-pituitary-adrenal (HPA) axis during periods of increased cytokine and/or neuropeptide productivity, such as during immune stimulation (Uribe et al., 1999;




Fig. 4. Effects of pretreatment with vehicle, L-NAME (30 mg/kg) or aminoguanidine (AG, 50 mg/kg) on food intake (A), and body weight change (B) evaluated at 24 h after administration of either LPS or saline. Each column represents the mean with S.E.M. The symbols denote significance levels: * $p < 0.05$; when compared with the control groups; # $p < 0.05$; when compared with the vehicle plus LPS group.

Bugajski et al., 2006; Gadek-Michalska and Bugajski, 2008; Jankord et al., 2009). Glucocorticoids have been suggested to play an important role in organisms coping with unpredictable events. During such events, glucocorticoids function to maintain allostasis or physiological stability (McEwen and Wingfield, 2003; Goymann and Wingfield, 2004; Romero et al., 2009). In our study, animals treated with nitric oxide synthase inhibitors prior to endotoxin administration displayed an increase in corticosterone secretion and accentuated depression-like behavior, anxiety and hypophagia. This finding suggests that NO is involved in the development of sickness behavior induced by LPS.

In conclusion, using a mouse model for acute systemic inflammation, we have shown that pharmacologically blocking NOS results in

accentuated LPS-induced behavioral changes. In addition, the present study suggests that NO may have a protective role, acting in an inhibitory feedback manner, to limit LPS-induced sickness behavior.

Disclosures

The authors indicate no potential conflicts of interest.

Acknowledgments

This work was supported by FAPEMIG, CNPq and CAPES. We are grateful for the excellent technical support of Marina F Venâncio and José dos Reis Pereira.




Fig. 5. Effects of pretreatment with vehicle, L-NAME (30 mg/kg) or aminoguanidine (AG, 50 mg/kg) on corticosterone plasma levels after administration of either LPS or saline. Each column represents the mean with S.E.M. The symbols denote significance levels: * $p < 0.05$; *** $p < 0.001$ when compared with the control groups; # $p < 0.05$; ### $p < 0.001$ when compared with the vehicle plus LPS group.

References

- Benedetti M, Merino R, Kusuda R, Ravanelli MI, Cadetti F, Dos Santos P, et al. Plasma corticosterone levels in mouse models of pain. *Eur J Pain* 2011. <http://dx.doi.org/10.1002/j.1532-2149.2011.00066.x>.
- Boyle III WA, Parvathaneni LS, Bourlier V, Sauter C, Laubach VE, Cobb JP. iNOS gene expression modulates microvascular responsiveness in endotoxin-challenged mice. *Circ Res* 2000;87:E18–24.
- Bugajski AJ, Gadek-Michalska A, Bugajski J. The involvement of nitric oxide and prostaglandins in the cholinergic stimulation of hypothalamic-pituitary-adrenal response during crowding stress. *J Physiol Pharmacol* 2006;57:463–77.
- Cauwels A. Nitric oxide in shock. *Kidney Int* 2007;72:557–65.
- Dallaporta M, Pecchi E, Jacques C, Berenbaum F, Jean A, Thirion S, et al. c-Fos immunoreactivity induced by intraperitoneal LPS administration is reduced in the brain of mice lacking the microsomal prostaglandin G synthase-1 (mPGES-1). *Brain Behav Immun* 2007;21:1109–21.
- Dantzer R. Cytokine, sickness behavior, and depression. *Immunol Allergy Clin North Am* 2009;29:247–64.
- Dantzer R, O'Connor JC, Freund GG, Johnson RW, Kelley KW. From inflammation to sickness and depression: when the immune system subjugates the brain. *Nat Rev Neurosci* 2008;9:46–56.
- de Paiva VN, Lima SN, Fernandes MM, Soncini R, Andrade CA, Giusti-Paiva A. Prostaglandins mediate depressive-like behaviour induced by endotoxin in mice. *Behav Brain Res* 2010;215:146–51.
- Dunn AJ, Swiergiel AH. Effects of interleukin-1 and endotoxin in the forced swim and tail suspension tests in mice. *Pharmacol Biochem Behav* 2005;81:688–93.
- Engblom D, Ek M, Saha S, Ericsson-Dahlstrand A, Jakobsson PJ, Blomqvist A. Prostaglandins as inflammatory messengers across the blood–brain barrier. *J Mol Med* 2002;80:5–15.
- Engblom D, Saha S, Engstrom L, Westman M, Audoly LP, Jakobsson PJ, et al. Microsomal prostaglandin G synthase-1 is the central switch during immune-induced pyresis. *Nat Neurosci* 2003;6:1137–8.

- Frenois F, Moreau M, O'Connor J, Lawson M, Micon C, Lestage J, et al. Lipopolysaccharide induces delayed FosB/DeltaFosB immunostaining within the mouse extended amygdala, hippocampus and hypothalamus, that parallel the expression of depressive-like behavior. *Psychoneuroendocrinology* 2007;32:516–31.
- Gadek-Michalska A, Bugajski J. Nitric oxide in the adrenergic-and CRH-induced activation of hypothalamic-pituitary-adrenal axis. *J Physiol Pharmacol* 2008;59:365–78.
- Giusti-Paiva A, De Castro M, Antunes-Rodrigues J, Carnio EC. Inducible nitric oxide synthase pathway in the central nervous system and vasopressin release during experimental septic shock. *Crit Care Med* 2002;30:1306–10.
- Giusti-Paiva A, Martinez MR, Felix JV, da Rocha MJ, Carnio EC, Elias LL, et al. Simvastatin decreases nitric oxide overproduction and reverts the impaired vascular responsiveness induced by endotoxemic shock in rats. *Shock* 2004;21:271–5.
- Giusti-Paiva A, Elias LL, Antunes-Rodrigues J. Inhibitory effect of gaseous neuromodulators in vasopressin and oxytocin release induced by endotoxin in rats. *Neurosci Lett* 2005;381:320–4.
- Goymann W, Wingfield JC. Allostatic load, social status and stress hormones: the costs of social status matter. *Anim Behav* 2004;67:591–602.
- Guidot DM, Repine MJ, Hybertson BM, Repine JE. Inhaled nitric oxide prevents neutrophil-mediated, oxygen radical-dependent leak in isolated rat lungs. *Am J Physiol* 1995;269:L2–5.
- Harada S, Imaki T, Chikada N. Distinct distribution and time-course changes in neuronal nitric oxide synthase and inducible NOS in the paraventricular nucleus following lipopolysaccharide injection. *Brain Res* 1999;821:322–32.
- Jankord R, McAllister RM, Ganjam VK, Laughlin MH. Chronic inhibition of nitric oxide synthase augments the ACTH response to exercise. *Am J Physiol Regul Integr Comp Physiol* 2009;296:R728–34.
- Kelley KW, Bluthé RM, Dantzer R, Zhou JH, Shen WH, Johnson RW, et al. Cytokine-induced sickness behavior. *Brain Behav Immun* 2003;17:112–8.
- Lacosta S, Merali Z, Anisman H. Behavioral and neurochemical consequences of lipopolysaccharide in mice: anxiogenic-like effects. *Brain Res* 1999;818:291–303.
- Long NC, Otterness I, Kunkel SL, Vander AJ, Kluger MJ. Roles of interleukin 1 beta and tumor necrosis factor in lipopolysaccharide fever in rats. *Am J Physiol* 1990;259:R724–8.
- Mahar Doan KM, Lakhman SS, Boje KM. Blood–brain barrier transport studies of organic guanidino cations using an in situ brain perfusion technique. *Brain Res* 2000;876:141–7.
- McEwen BS, Wingfield JC. The concept of allostasis in biology and biomedicine. *Horm Behav* 2003;43:2–15.
- Metcalfe K, Jungersten L, Lisander B. Effective inhibition of nitric oxide production by aminoguanidine does not reverse hypotension in endotoxaemic rats. *Acta Anaesthesiol Scand* 2002;46:17–23.
- Minc-Golomb D, Tsarfaty I, Schwartz JP. Expression of inducible nitric oxide synthase by neurones following exposure to endotoxin and cytokine. *Br J Pharmacol* 1994;112:720–2.
- Mollace V, Muscoli C, Masini E, Cuzzocrea S, Salvemini D. Modulation of prostaglandin biosynthesis by nitric oxide and nitric oxide donors. *Pharmacol Rev* 2005;57:217–52.
- Moncada S, Palmer RM, Higgs EA. Nitric oxide: physiology, pathophysiology and pharmacology. *Pharmacol Rev* 1991;43:109–42.
- Parnet P, Kelley KW, Bluthé RM, Dantzer R. Expression and regulation of interleukin-1 receptors in the brain. Role in cytokines-induced sickness behavior. *J Neuroimmunol* 2002;125:5–14.
- Porsolt RD, Bertin A, Jalfre M. Behavioral despair in mice: a primary screening test for antidepressants. *Arch Int Pharmacodyn Ther* 1977;229:327–36.
- Romero LM, Dickens MJ, Cyr NE. The reactive scope model – a new model integrating homeostasis, allostasis, and stress. *Horm Behav* 2009;55:375–89.
- Rorato R, Menezes AM, Giusti-Paiva A, de Castro M, Antunes-Rodrigues J, Elias LL. Prostaglandin mediates endotoxaemia-induced hypophagia by activation of pro-opiomelanocortin and corticotrophin-releasing factor neurons in rats. *Exp Physiol* 2009;94:371–9.
- Soncini R, de Souza DF, Neves AP, Braga DS, Andrade CA, Giusti-Paiva A. Dipyrone attenuates acute sickness response to lipopolysaccharide in mice. *Neurosci Lett* 2012;516:114–8.
- Szabo C. Physiological and pathophysiological roles of nitric oxide in the central nervous system. *Brain Res Bull* 1996;41:131–41.
- Thiemermann C. Nitric oxide and septic shock. *Gen Pharmacol* 1997;29:159–66.
- Trzeciak S, Cinel I, Phillip Dellinger R, Shapiro NI, Arnold RC, Parrillo JE, et al. Microcirculatory Alterations in Resuscitation and Shock (MARS) Investigators. Resuscitating the microcirculation in sepsis: the central role of nitric oxide, emerging concepts for novel therapies, and challenges for clinical trials. *Acad Emerg Med* 2008;15:399–413.
- Turnbull AV, Rivier CL. Regulation of the hypothalamic-pituitary-adrenal axis by cytokines: actions and mechanisms of action. *Physiol Rev* 1999;79:1–71.
- Uribe RM, Lee S, Rivier C. Endotoxin stimulates nitric oxide production in the paraventricular nucleus of the hypothalamus through nitric oxide synthase I: correlation with hypothalamic-pituitary-adrenal axis activation. *Endocrinology* 1999;140:5971–81.
- Ushikubi F, Segi E, Sugimoto Y, Murata T, Matsuoka T, Kobayashi T, et al. Impaired febrile response in mice lacking the prostaglandin E receptor subtype EP3. *Nature* 1998;395:281–4.
- Vincent JL, Zhang H, Szabo C, Freiser JH. Effects of nitric oxide in septic shock. *Am J Respir Crit Care Med* 2000;161:1781–5.
- Vona-Davis L, Wearden P, Hill J, Hill R. Cardiac response to nitric oxide synthase inhibition using aminoguanidine in a rat model of endotoxemia. *Shock* 2002;17:404–10.